

Trying 9158046...Open

box200> enter system id

Logging in to Dialog

DIALOG INFORMATION SERVICES

PLEASE LOGON:

\*\*\*\*\*

IALOG Invalid account number

DIALOG INFORMATION SERVICES

PLEASE LOGON:

\*\*\*\*\*

ENTER PASSWORD:

717093fe

\*\*\*\*\*

Welcome to DIALOG

Dialog level 98.07.06D

Reconnected in file OS 06aug98 16:49:30

\* \* \* NEW RATES STRUCTURE

\* \* \* Effective June 1, connect time charges on Dialog have been

\* \* \* eliminated and DialUnits charges have been introduced.

\* \* \* Please check HomeBase for the text of the press release

\* \* \* announcing this change.

\* \* \*

\* \* \* The ERIC Dialorder supplier now requires prepayment with

\* \* \* all orders. For information contact ERIC document supply

\* \* \* at 800-443-3742 or service@edrs.com.

\* \* \*

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1966-1998/Sep W4

(c) format only 1998 Dialog Corporation

File 5:BIOSIS PREVIEWS(R) 1969-1998/JUL W4

(c) 1998 BIOSIS

File 73:EMBASE 1974-1998/Aug W2

(c) 1998 Elsevier Science B.V.

File 144:Pascal 1973-1998/Jul

(c) 1998 INIST/CNRS

File 434:SciSearch(R) Cited Ref Sci 1974-1989/Dec

(c) 1998 Inst for Sci Info

File 34:SciSearch(R) Cited Ref Sci 1990-1998/Jul W4

(c) 1998 Inst for Sci Info

File 76:Life Sciences Collection 1982-1998/Jun

(c) 1998 Cambridge Sci Abs

File 305:Analytical Abstracts 1980-1998/Aug

(c) 1998 Royal Soc Chemistry

File 156:Toxline(R) 1965-1998/Jul

(c) format only 1998 The Dialog Corporation

File 442:AMA Journals 1982-1998/Jul W4

(c) 1998 Amer Med Assn -FARS/DARS apply

File 35:Dissertation Abstracts Online 1861-1998/Aug

(c) 1998 UMI

File 50:CAB Abstracts 1972-1998/Jun  
 (c) 1998 CAB International  
 File 77:Conference Papers Index 1973-1998/Jul  
 (c) 1998 Cambridge Sci Abs  
 File 150:Legal Resource Index(TM) 1980-1998/Aug 06  
 (c) 1998 Info. Access Co.  
 File 376:Derwent Drug File 1964-1982  
 (c) 1995 Derwent Info Ltd.  
 File 2:INSPEC 1969-1998/Aug W2  
 (c) 1998 Institution of Electrical Engineers  
 File 8:Ei Compendex(R) 1970-1998/Aug W4  
 (c) 1998 Engineering Info. Inc.  
 File 103:Energy SciTec 1974-1998/Jul B1  
 (c) 1998 Contains copyrighted material  
 \*File 103: For access restrictions, see HELP RESTRICT.  
 File 348:EUROPEAN PATENTS 1978-1998/Jul W3-  
 (c) 1998 EUROPEAN PATENT OFFICE  
 \*File 348: \*\* NEW FEATURE \*\* English language translations of French  
 and German abstracts now searchable. See HELP NEWS 348 for info.  
 File 653:US Patents Fulltext 1980-1989  
 (c) format only 1998 The Dialog Corp.  
 \*File 653: Reassignment data now current through 05/14/98.  
 Reexamination, extension, expiration, reinstatement updated weekly.  
 File 654:US Pat.Full. 1990-1998/Aug 04  
 (c) format only 1998 The Dialog Corp.  
 \*File 654: Reassignment data now current through 05/14/98.  
 Reexamination, extension, expiration, reinstatement updated weekly.

Set	Items	Description
?	set hi	;set hi

Hilight option is not available in file(s) 77  
 HILIGHT set on as ''  
**Hilight option is not available in file(s) 77**  
**HILIGHT set on as ''**  
 ? ds

Set	Items	Description
S1	366	(ERYTHROCYT? OR RED) (4N) (ADENYLATE(W) KINASE)
S2	227	RD (unique items)
? t s2/7/2,3,20,24,32,50,91,104,145,165,171,174		

2/7/2 (Item 2 from file: 155)  
 DIALOG(R)File 155:MEDLINE(R)  
 (c) format only 1998 Dialog Corporation. All rts. reserv.

09371773 98069130

**Erythrocyte adenylate kinase** isoenzyme as a marker for hemolysis.

Thomas G; Murthy VV  
 Department of Pathology, Albert Einstein College of Medicine, Bronx, New York, USA.

J Clin Lab Anal (UNITED STATES) 1997, 11 (6) p351-6, ISSN 0887-8013  
 Journal Code: JLA

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The presence in serum of **adenylate kinase** isoenzyme

originating from **erythrocyte** can be useful as a marker for detecting hemolysis. We have presented preliminary evidence for identifying hemolytic anemia patients earlier by determining erythrocyte AK isoenzyme activity in serum (or plasma) rather than using measurement of plasma hemoglobin concentration. This test being quite specific for hemolysis should find use as a quick method for estimating the extent of in vivo hemolysis in hemolytic patients earlier than heretofore possible.

2/7/3 (Item 3 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

09138555 97362603

Differentiation and resolution of **erythrocyte** and muscle **adenylate kinase** activities in serum by electrophoresis.  
Murthy VV; Ali F; Burns ER  
Department of Pathology, Albert Einstein College of Medicine, The Bronx, New York, USA.

J Clin Lab Anal (UNITED STATES) 1997, 11 (4) p235-7, ISSN 0887-8013  
Journal Code: JLA

Languages: ENGLISH

Document type: JOURNAL ARTICLE

**Adenylate kinase** activity originating from **erythrocytes** has been shown to be distinct from muscle adenylate kinase or myokinase activity, until now considered to be identical enzyme activities. The two activities can be differentiated by electrophoretic fractionation, thus making it possible to quantify the **erythrocyte adenylate kinase** activity present in serum.

2/7/20 (Item 20 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

05530841 89133735

The effect of hemolysis on creatine kinase determination [see comments]  
Greenson JK; Farber SJ; Dubin SB  
Department of Pathology and Laboratory Medicine, Cedars-Sinai Medical Center, Los Angeles, CA 90048.

Arch Pathol Lab Med (UNITED STATES) Feb 1989, 113 (2) p184-5, ISSN 0003-9985 Journal Code: 79Z

Comment in Arch Pathol Lab Med 1992 Jan;116(1):7-8

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Hemolysis can cause falsely elevated creatine kinase (CK) values when spectrophotometric methods of measurement are used. This apparent increase in CK is due to the **red** blood cell enzyme **adenylate kinase**. In an attempt to reduce this interference, most commercial CK kits employ adenosine monophosphate and/or diadenosine pentaphosphate as adenylate kinase inhibitors. To determine whether hemolyzed specimens should be accepted for testing, we measured the CK values of 26 serum samples, each with six different concentrations of added hemolysate. The results showed that hemolysis had an additive effect on CK, with an average increase in CK of approximately 10 U/L for every 1 g/L of hemoglobin. In most settings, this increase is not clinically significant. In the case of massive hemolysis, the hemoglobin concentration of the serum can be measured to correct the apparent CK value. The exclusion of hemolyzed specimens is unnecessary.

2/7/24 (Item 24 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

04427447 84289246  
Leakage of adenylate kinase from stored blood cells.  
Olsson T; Gulliksson H; Palmeborn M; Bergstrom K; Thore A  
J Appl Biochem (UNITED STATES) Dec 1983, 5 (6) p437-45, ISSN  
0161-7354 Journal Code: HEA  
Languages: ENGLISH  
Document type: JOURNAL ARTICLE

The bioluminescent firefly luciferase assay for ATP was used to measure adenylate kinase activity in plasma. The formation of ATP from ADP was measured continuously in a coupled assay using a luminometer. Optimal analytical conditions were determined for the coupled reaction. The assay was used to follow accumulation of adenylate kinase in plasma of different preparations of stored **red** blood cells. **Adenylate kinase** was found to be released concomitantly with hemoglobin during aging. There was a high degree of correlation between the amount of accumulated hemoglobin and adenylate kinase. The assay was also used to measure lysis of stored platelets during aging.

2/7/32 (Item 32 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

04307438 81134512  
The primary cause of hemolysis in enzymopathies of anaerobic glycolysis: a viewpoint.  
Valentine WN; Paglia DE  
Blood Cells (GERMANY, WEST) 1980, 6 (4) p819-29, ISSN 0340-4684  
Journal Code: A8H  
Languages: ENGLISH  
Document type: JOURNAL ARTICLE

2/7/50 (Item 50 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

03055042 79064127  
Anion-exchange chromatography of **erythrocytic** and muscle **adenylate kinase** and its effect on the serum creatine kinase isoenzyme assays.  
Klein B; Jeunelot CL  
Clin Chem (UNITED STATES) Dec 1978, 24 (12) p2168-70, ISSN 0009-9147  
Journal Code: DBZ  
Languages: ENGLISH  
Document type: JOURNAL ARTICLE  
We determined the elution profile of erythrocytic and muscle adenylate kinases (EC 2.7.4.3) in the Roche chromatographic creatine kinase procedure and studied the interference these enzymes would cause in the isolation and assay of serum creatine kinase (EC 2.7.3.2) isoenzymes. Both adenylate kinases co-elute with the creatine kinase MM fraction and do not interfere with the isolation or assay of the MB fraction.

2/7/91 (Item 3 from file: 5)

DIALOG(R)File 5:BIOSIS PREVIEWS(R)  
(c) 1998 BIOSIS. All rts. reserv.

13354942 BIOSIS Number: 99354942

Clinical utility of serum **erythrocyte adenylate kinase**,  
a new marker for hemolysis

Kale A; Murthy V V; Burns E R

Dep. Pathol., Albert Einstein Coll. Med., Bronx, NY, USA

Blood 88 (10 SUPPL. 1 PART 1-2). 1996. 5B.

Full Journal Title: Thirty-eighth Annual Meeting of the American Society  
of Hematology, Orlando, Florida, USA, December 6-10, 1996. Blood

ISSN: 0006-4971

Language: ENGLISH

Print Number: Biological Abstracts/RRM Vol. 049 Iss. 002 Ref. 027834

2/7/104 (Item 16 from file: 5)  
DIALOG(R)File 5:BIOSIS PREVIEWS(R)  
(c) 1998 BIOSIS. All rts. reserv.

4210680 BIOSIS Number: 26063023

**ADENYLATE KINASE FROM HUMAN ERYTHROCYTES AND PLATELETS**

NEALON D A; PRIDGAR E; HENDERSON A R

CTR. LABS. RES., NY STATE DEP. HEALTH, ALBANY, NY 12201.

34TH NATIONAL MEETING OF THE AMERICAN ASSOCIATION FOR CLINICAL CHEMISTRY,  
ANAHEIM, CALIF., USA, AUG. 8-13, 1982. CLIN CHEM 28 (7). 1982. 1606.

CODEN: CLCHA

Language: ENGLISH

2/7/145 (Item 2 from file: 73)  
DIALOG(R)File 73:EMBASE  
(c) 1998 Elsevier Science B.V. All rts. reserv.

10601611 EMBASE No: 98029364

Effects of haemolysis on the Boehringer Mannheim creatine kinase-MB assay  
Donnelly J.G.

J.G. Donnelly, Department of Laboratory Medicine, Ottawa Civic Hospital,  
University of Ottawa School Medicine, 1053 Carling Avenue, Ottawa, Ont. K1Y  
4E9 Canada

Annals of Clinical Biochemistry (United Kingdom) , 1998, 35/1 (143-144)

CODEN: ACBOB ISSN: 0004-5632

DOCUMENT TYPE: Journal Article

LANGUAGES: ENGLISH SUMMARY LANGUAGES: ENGLISH

NUMBER OF REFERENCES: 3

The Boehringer Mannheim (BM; Boehringer Mannheim, Laval Quebec, Canada)  
assay for creatine kinase-B (CK-MB) measures the residual catalytic  
activity of the creatine kinase-B subunit after immunoinhibition of the M  
subunit. During our evaluation of this assay for implementation on the  
Hitachi 917 analyser we observed a profound positive interference from  
haemolysis. While the effect of **erythrocyte adenylate  
kinase** is widely known to users of CK assays, we found that there was  
significant interference even when the haemolysis could not be visually  
detected. We investigated the extent of this interference in order to  
determine the suitability of haemolysed specimens for this assay.

2/7/165 (Item 22 from file: 73)  
DIALOG(R)File 73:EMBASE  
(c) 1998 Elsevier Science B.V. All rts. reserv.

817455 EMBASE No: 77200490

Agarose thin layer electrophoresis for the determination of red  
cell **adenylate kinase** (EC 2.7.4.3) polymorphisms  
AGAROSE DUNNSCHICHT ELEKTROPHORESE ZUR BESTIMMUNG DER ERYTHROZYTAREN  
ADENYLATKINASE (EC 2.7.4.3) POLYMORPHISMEN

Tsuji T.; Weissmann J.

Abt. Rechtsmed., Med. Hochsch., Lubeck GERMANY, WEST

ARZTL.LAB. (GERMANY, WEST) , 1976, 22/11 (363-365)

CODEN: AELAA

LANGUAGES: GERMAN

A simple method for the determination of AK phenotypes by means of  
agarose thin layer electrophoresis is reported and compared with the agar  
and CAM methods. Separation was excellent and the spots were well  
demarcated. The results were better than those obtained with the other two  
methods.

2/7/171 (Item 28 from file: 73)

DIALOG(R)File 73:EMBASE

(c) 1998 Elsevier Science B.V. All rts. reserv.

462506 EMBASE No: 76043888

Studies on storage induced changes of isoenzyme patterns (PGM, AK, ADA)  
by means of cellulose acetate membrane (CAM) electrophoresis

UNTERSUCHUNGEN UBER LAGERUNGSBEDINGTE VERANDERUNGEN VON ISOENZYMUSTERN  
(PGM, AK, ADA) MIT HILFE DER CAF ELEKTROPHORESE

Berndt H.; Kox N.

Abt. Immunol. Transf. Med., Med. Hochsch., Lubeck GERMANY, WEST

ARZTL.LAB. (GERMANY, WEST) , 1975, 21/3 (87-97)

CODEN: AELAA

LANGUAGES: GERMAN

2/7/174 (Item 31 from file: 73)

DIALOG(R)File 73:EMBASE

(c) 1998 Elsevier Science B.V. All rts. reserv.

290293 EMBASE No: 75081287

Isolation and properties of isoenzymes of adenylate kinase  
REINIGUNG UND EIGENSCHAFTEN VON ISOENZYMEN DER ADENYLATKINASE

Mebis D.

Zent. Rechtsmed., Univ. Frankfurt/M. GERMANY, WEST

BEITR.GERICHTL.MED. (--) , 1973, No:31 (295-296)

CODEN: BEGMA

LANGUAGES: GERMAN

The isoenzymes of adenylate kinase were isolated from the hemolysate of  
pig erythrocytes by ammonium sulfate fractionation, pH treatment, gel  
filtration and ion exchange chromatography. The enzyme specifically  
catalyzes the reaction 2 adenosine diphosphate < - > adenosine triphosphate  
+ adenosine monophosphate. Its molecular weight was found to be 23,500 by  
gel filtration.

? t s2/pn/226

>>>No matching display code(s) found in file(s): 8, 34, 50, 73, 155-156,  
305, 376, 434, 442

2/PN/226 (Item 1 from file: 653)

DIALOG(R)File 653:(c) format only 1998 The Dialog Corp. All rts. reserv.

PATENT NO.: 4,220,714  
ISSUED: September 02, 1980 (19800902)  
? s s2 and (hemolysis or hemolyzed)

227 S2  
68804 HEMOLYSIS  
2273 HEMOLYZED  
S3 16 S2 AND (HEMOLYSIS OR HEMOLYZED)  
? t s3/7/1-16

3/7/1 (Item 1 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

09371773 98069130

**Erythrocyte adenylate kinase** isoenzyme as a marker for  
**hemolysis**.

Thomas G; Murthy VV  
Department of Pathology, Albert Einstein College of Medicine, Bronx, New  
York, USA.

J Clin Lab Anal (UNITED STATES) 1997, 11 (6) p351-6, ISSN 0887-8013

Journal Code: JLA

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The presence in serum of **adenylate kinase** isoenzyme  
originating from **erythrocyte** can be useful as a marker for detecting  
**hemolysis**. We have presented preliminary evidence for identifying  
hemolytic anemia patients earlier by determining erythrocyte AK isoenzyme  
activity in serum (or plasma) rather than using measurement of plasma  
hemoglobin concentration. This test being quite specific for  
**hemolysis** should find use as a quick method for estimating the extent  
of in vivo **hemolysis** in hemolytic patients earlier than heretofore  
possible.

3/7/2 (Item 2 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

09138555 97362603

Differentiation and resolution of **erythrocyte** and muscle  
**adenylate kinase** activities in serum by electrophoresis.

Murthy VV; Ali F; Burns ER  
Department of Pathology, Albert Einstein College of Medicine, The Bronx,  
New York, USA.

J Clin Lab Anal (UNITED STATES) 1997, 11 (4) p235-7, ISSN 0887-8013

Journal Code: JLA

Languages: ENGLISH

Document type: JOURNAL ARTICLE

**Adenylate kinase** activity originating from **erythrocytes**  
has been shown to be distinct from muscle adenylate kinase or myokinase  
activity, until now considered to be identical enzyme activities. The two  
activities can be differentiated by electrophoretic fractionation, thus  
making it possible to quantify the **erythrocyte adenylate**  
**kinase** activity present in serum.

3/7/3 (Item 3 from file: 155)  
DIALOG(R)File 155:MEDLINE(R)

(c) format only 1998 Dialog Corporation. All rts. reserv.

08795088 97045577

[Hemolytic anemia due to abnormalities in erythrocyte nucleotide metabolism]

Masuda M; Mizoguchi H

Department of Hematology, Tokyo Women's Medical College.

Nippon Rinsho (JAPAN) Sep 1996, 54 (9) p2473-7, ISSN 0047-1852

Journal Code: KIM

Languages: JAPANESE Summary Languages: ENGLISH

Document type: JOURNAL ARTICLE; REVIEW; REVIEW LITERATURE English  
Abstract

Abnormalities in erythrocyte nucleotide metabolism are associated with hereditary nonspherocytic hemolytic anemia. Deficiency of adenylate kinase and pyrimidine 5'-nucleotidase and hyperactivity of adenosine deaminase shorten the **red** cell lifespan. Deficiency of **adenylate kinase** has been reported in four different families. Although in one family, total absence of this enzymatic activity was documented in one hematologically normal sibling, there is doubt about the capacity of this single enzyme deficiency to produce **hemolysis**. A deficiency of pyrimidine 5'-nucleotidase is a cause of hemolytic anemia characterized by red cells with basophilic stippling. This enzyme has been reported to catalyze the hydrolytic dephosphorylation of pyrimidine 5'-ribose monophosphate. Red cells of patients contain an increased concentration of pyrimidine nucleotides and reduced form of glutathione. In hyperactivity, the adenosine deaminase activity in erythrocytes may be increased to 100 times the normal level. The high adenosine deaminase activity of erythrocytes depletes adenine nucleotides, inhibiting its metabolism. (15 Refs.)

3/7/4 (Item 4 from file: 155)

DIALOG(R) File 155: MEDLINE(R)

(c) format only 1998 Dialog Corporation. All rts. reserv.

08224518 94322148

Adenylate kinase mimics creatine kinase-MM isoenzyme in a CK isoenzyme electrophoresis assay.

Murthy VV

Department of Laboratory Medicine, Albert Einstein College of Medicine, Bronx, New York.

J Clin Lab Anal (UNITED STATES) 1994, 8 (3) p140-3, ISSN 0887-8013

Journal Code: JLA

Languages: ENGLISH

Document type: JOURNAL ARTICLE

**Adenylate kinase** activity (AK) originating from **erythrocytes**, present in **hemolyzed** serum behaves like creatine kinase MM isoenzyme (CK-MM) in some CK electrophoresis assays that employ, in their visualization reagent kits, adenosine monophosphate (AMP) as the sole inhibitor of AK, rather than a combination of AMP and a more potent inhibitor of erythrocyte AK, diadenosine pentaphosphate (Ap5A), to inhibit all contaminating-AK activities in serum and quantify only the CK isoenzyme activities in serum following electrophoretic fractionation on agarose gel. This can spuriously overestimate the CK-MM fraction and thereby result in underestimation of CK-MM or CK-BB isoenzymes if present. A **hemolyzed** serum sample obtained from an elderly patient was erroneously reported as containing low CK-MM due to such overestimation of CK-MM fraction in the sample. Supplementing the AMP already present in the visualization reagent formulation, used to estimate CK isoenzyme concentration in serum, with Ap5A can eliminate or effectively minimize AK interference, especially that



caused by **hemolysis** , and thereby prevent reporting false-negative CK-MB result obtained with CK isoenzyme electrophoresis assays.

3/7/5 (Item 5 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

05530841 89133735

The effect of **hemolysis** on creatine kinase determination [see comments]

Greenson JK; Farber SJ; Dubin SB  
Department of Pathology and Laboratory Medicine, Cedars-Sinai Medical Center, Los Angeles, CA 90048.

Arch Pathol Lab Med (UNITED STATES) Feb 1989, 113 (2) p184-5, ISSN 0003-9985 Journal Code: 79Z

Comment in Arch Pathol Lab Med 1992 Jan;116(1):7-8

Languages: ENGLISH

Document type: JOURNAL ARTICLE

**Hemolysis** can cause falsely elevated creatine kinase (CK) values when spectrophotometric methods of measurement are used. This apparent increase in CK is due to the **red** blood cell enzyme **adenylate kinase**. In an attempt to reduce this interference, most commercial CK kits employ adenosine monophosphate and/or diadenosine pentaphosphate as adenylylase kinase inhibitors. To determine whether **hemolyzed** specimens should be accepted for testing, we measured the CK values of 26 serum samples, each with six different concentrations of added hemolysate. The results showed that **hemolysis** had an additive effect on CK, with an average increase in CK of approximately 10 U/L for every 1 g/L of hemoglobin. In most settings, this increase is not clinically significant. In the case of massive **hemolysis**, the hemoglobin concentration of the serum can be measured to correct the apparent CK value. The exclusion of **hemolyzed** specimens is unnecessary.

3/7/6 (Item 6 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

04404283 83266230

Metabolic compensation for profound **erythrocyte adenylylase kinase** deficiency. A hereditary enzyme defect without hemolytic anemia.

Beutler E; Carson D; Dannawi H; Forman L; Kuhl W; West C; Westwood B  
J Clin Invest (UNITED STATES) Aug 1983, 72 (2) p648-55, ISSN 0021-9738 Journal Code: HS7

Contract/Grant No.: HL 25552, HL, NHLBI

Languages: ENGLISH

Document type: JOURNAL ARTICLE

A child with hemolytic anemia was found to have severe **erythrocyte adenylylase kinase** (AK) deficiency, but an equally enzyme-deficient sibling had no evidence of **hemolysis**. No residual enzyme activity was found in erythrocytes by spectrophotometric methods that could easily have detected 0.1% of normal activity. However, concentrated hemolysates were shown to have the capacity to generate small amounts of ATP and AMP from ADP after prolonged incubation. Hemolysates could also catalyze the transfer of labeled gamma-phosphate from ATP to ADP. Intact erythrocytes were able to transfer phosphate from the gamma-position of ATP to the beta-position, albeit at a rate substantially slower than normal. They could also incorporate <sup>14</sup>C-labeled adenine into

ADP and ATP. Thus, a small amount of residual AK-like activity representing about 1/2,000 of the activity normally present could be documented in the deficient erythrocytes. The residual activity was not inhibited by N-ethylmaleimide, which completely abolishes the activity of the normal AK1 isozyme of erythrocytes. The minute amount of residual activity in erythrocytes could represent a small amount of the AK2 isozyme, which has not been thought to be present in erythrocytes, or the activity of erythrocyte guanylate kinase with AMP substituting as substrate for GMP. Peripheral blood leukocytes, cultured skin fibroblasts, and transformed lymphoblasts from the deficient subject manifested about 17, 24, and 74%, respectively, of the activity of the concurrent controls. This residual activity is consistent with the existence of genetically independent AK isozyme, AK2, which is known to exist in these tissues. The cause of **hemolysis** in the proband was not identified. Possibilities include an unrelated enzyme deficiency or other erythrocyte enzyme defect and intraction of another unidentified defect with AK deficiency.

3/7/7 (Item 7 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

04401727 83228082

**Red cell adenylate kinase** deficiency associated with hereditary nonspherocytic hemolytic anemia: clinical and biochemical studies.

Miwa S; Fujii H; Tani K; Takahashi K; Takizawa T; Igarashi T  
Am J Hematol (UNITED STATES) Jun 1983, 14 (4) p325-33, ISSN 0361-8609  
Journal Code: 3H4

Languages: ENGLISH

Document type: JOURNAL ARTICLE

We report here a case of **red cell adenylate kinase (AK)** deficiency associated with hereditary hemolytic anemia. The proband is a 10-year-old Japanese girl. Her physical and mental development was normal. She has shown moderate to mild hemolytic anemia since the neonatal period and hepatosplenomegaly. The red cell AK activity was 44% of normal. Contents of red cell glycolytic intermediates and adenine nucleotides were normal when compared with a comparable reticulocyte-rich control. Glucose consumption and lactate formation were normal. Hexose monophosphate shunt activity was somewhat lower than that of a comparable reticulocyte-rich control. There were no significant differences in the contents of adenine nucleotides between the younger and older red cells of the patient. Enzymatic characterization by hemolysate revealed that the patient's AK had an increased Michaelis constant for adenosine diphosphate and slight thermal instability. The patient's enzyme migrated approximately half-way between the AK 1 and AK 2 position on starch-gel electrophoresis. The mode of inheritance of this case is obscure. The mechanism of **hemolysis** might be a structural gene mutation that caused altered electrophoretic and kinetic properties.

3/7/8 (Item 8 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

04307438 81134512

The primary cause of **hemolysis** in enzymopathies of anaerobic glycolysis: a viewpoint.

Valentine WN; Paglia DE

Blood Cells (GERMANY, WEST) 1980, 6 (4) p819-29, ISSN 0340-4684

Journal Code: A8H  
Languages: ENGLISH  
Document type: JOURNAL ARTICLE

3/7/9 (Item 9 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

03325189 81013018  
Increased creatine kinase activities associated with haemolysis.  
Bais R; Edwards JB  
Pathology (AUSTRALIA) Apr 1980, 12 (2) p203-7, ISSN 0031-3025  
Journal Code: OTA  
Languages: ENGLISH  
Document type: JOURNAL ARTICLE  
The effect of haemolysis on creatine kinase activity has been investigated. The presence of **adenylate kinase** released from **erythrocytes** increases the apparent creatine kinase activity. This can be overcome by the addition of 10  $\mu\text{mol/l}$  of diadenosine pentaphosphate to the reagents.

3/7/10 (Item 10 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

03029062 77024503  
Creatine kinase in serum: 2. Interference of adenylate kinase with the assay.  
Szasz G; Gerhardt W; Gruber W; Bernt E  
Clin Chem (UNITED STATES) Nov 1976, 22 (11) p1806-11, ISSN 0009-9147  
Journal Code: DBZ  
Languages: ENGLISH  
Document type: JOURNAL ARTICLE  
Interference of adenylate kinase with Oliver's method [Biochem. J. 61, 116 (1955)] for creatine kinase is usually suppressed by including an adenylate kinase inhibitor, AMP. We studied the kinetics and compared the inhibition capacities of AMP and diadenosine pentaphosphate. Both are competitive inhibitors, AMP being markedly weaker, with a  $K_i$  of about 300  $\mu\text{mol/liter}$  for **adenylate kinase** from **erythrocyte**, muscle, and liver. AMP also weakly inhibits creatine kinase. Diadenosine pentaphosphate inhibits **erythrocyte** and muscle **adenylate kinase** strongly ( $K_i$  about 0.03  $\mu\text{mol/liter}$ ), the liver isoenzyme less strongly ( $K_i$  about 3  $\mu\text{mol/liter}$ ), and has no effect on creatine kinase up to 100  $\mu\text{mol/liter}$ . All three adenylate kinases may be present in a patient's serum, causing sample blanks to be high in a creatine kinase assay that lacks inhibitors. In acute hepatic damage, liver adenylate kinase activity in serum can be grossly increased. Use of sufficient diadenosine pentaphosphate alone for complete inhibition is relatively expensive. Consequently, we recommend a combination of both inhibitors. Diadenosine pentaphosphate, 10  $\mu\text{mol}$ , combined with 5  $\text{mmol}$  of AMP per liter inhibits **adenylate kinase** from **erythrocytes** and muscle by 97% and from liver by 95%.

3/7/11 (Item 11 from file: 155)  
DIALOG(R) File 155:MEDLINE(R)  
(c) format only 1998 Dialog Corporation. All rts. reserv.

00214938 68002942

The **adenylate kinase** of human plasma, **erythrocytes** and platelets in relation to the degradation of adenosine diphosphate in plasma.

Haslam RJ; Mills DC

Biochem J (ENGLAND) Jun 1967, 103 (3) p773-84, ISSN 0006-2936

Journal Code: 9YO

Languages: ENGLISH

Document type: JOURNAL ARTICLE

3/7/12 (Item 1 from file: 5)

DIALOG(R)File 5:BIOSIS PREVIEWS(R)

(c) 1998 BIOSIS. All rts. reserv.

13354942 BIOSIS Number: 99354942

Clinical utility of serum **erythrocyte adenylate kinase**, a new marker for **hemolysis**

Kale A; Murthy V V; Burns E R

Dep. Pathol., Albert Einstein Coll. Med., Bronx, NY, USA

Blood 88 (10 SUPPL. 1 PART 1-2). 1996. 5B.

Full Journal Title: Thirty-eighth Annual Meeting of the American Society of Hematology, Orlando, Florida, USA, December 6-10, 1996. Blood

ISSN: 0006-4971

Language: ENGLISH

Print Number: Biological Abstracts/RRM Vol. 049 Iss. 002 Ref. 027834

3/7/13 (Item 2 from file: 5)

DIALOG(R)File 5:BIOSIS PREVIEWS(R)

(c) 1998 BIOSIS. All rts. reserv.

3278089 BIOSIS Number: 71000488

INCREASED CREATINE KINASE EC-2.7.3.2 ACTIVITIES ASSOCIATED WITH **HEMOLYSIS**

BAIS R; EDWARDS J B

DIV. CLIN. CHEM., INST. MED. VET. SCI., FROME RD., ADELAIDE, S. AUST. 5000, AUST.

PATHOLOGY 12 (2). 1980. 203-207. CODEN: PTLGA

Full Journal Title: Pathology

Language: ENGLISH

The effect of **hemolysis** on creatine kinase [EC 2.7.3.2] activity was investigated. The presence of **adenylate kinase** released from **erythrocytes** increases the apparent creatine kinase activity. This can be overcome by the addition of 10  $\mu\text{mol/l}$  of diadenosine pentaphosphate to the reagents.

3/7/14 (Item 3 from file: 5)

DIALOG(R)File 5:BIOSIS PREVIEWS(R)

(c) 1998 BIOSIS. All rts. reserv.

3159871 BIOSIS Number: 20022278

EVALUATION OF NEW CREATINE KINASE FORMULATION ON ABBOTT BI CHROMATIC ANALYZERS

NERI B P; OLSON R M; ELSE R C

ABBOTT DIAGNOSTICS, N. CHICAGO, IL 60064.

JOINT MEETING OF THE AMERICAN ASSOCIATION FOR CLINICAL CHEMISTRY AND THE CANADIAN SOCIETY OF CLINICAL CHEMISTS, BOSTON, MASS., USA, JULY 20-25, 1980. CLIN CHEM 26 (7). 1980. 996-997. CODEN: CLCHA

Language: ENGLISH

3/7/15 (Item 1 from file: 73)  
DIALOG(R)File 73:EMBASE  
(c) 1998 Elsevier Science B.V. All rts. reserv.

10601611 EMBASE No: 98029364  
Effects of haemolysis on the Boehringer Mannheim creatine kinase-MB assay  
Donnelly J.G.  
J.G. Donnelly, Department of Laboratory Medicine, Ottawa Civic Hospital,  
University of Ottawa School Medicine, 1053 Carling Avenue, Ottawa, Ont. K1Y  
4E9 Canada  
Annals of Clinical Biochemistry (United Kingdom) , 1998, 35/1 (143-144)  
CODEN: ACBOB ISSN: 0004-5632  
DOCUMENT TYPE: Journal Article  
LANGUAGES: ENGLISH SUMMARY LANGUAGES: ENGLISH  
NUMBER OF REFERENCES: 3  
The Boehringer Mannheim (BM; Boehringer Mannheim, Laval Quebec, Canada)  
assay for creatine kinase-B (CK-MB) measures the residual catalytic  
activity of the creatine kinase-B subunit after immunoinhibition of the M  
subunit. During our evaluation of this assay for implementation on the  
Hitachi 917 analyser we observed a profound positive interference from  
haemolysis. While the effect of **erythrocyte adenylate**  
**kinase** is widely known to users of CK assays, we found that there was  
significant interference even when the haemolysis could not be visually  
detected. We investigated the extent of this interference in order to  
determine the suitability of haemolysed specimens for this assay.

3/7/16 (Item 2 from file: 73)  
DIALOG(R)File 73:EMBASE  
(c) 1998 Elsevier Science B.V. All rts. reserv.

957323 EMBASE No: 78125886  
Carrier detection in X linked recessive (Duchenne) muscular dystrophy:  
pyruvate kinase isoenzymes and creatine phosphokinase in serum and blood  
cells  
Smith I.; Thomson W.H.S.  
Res. Lab., Knightswood Hosp., Glasgow UNITED KINGDOM  
CLIN.CHIM.ACTA (NETHERLANDS) , 1977, 78/3 (439-451)  
CODEN: CCATA  
LANGUAGES: ENGLISH  
Allosterism allows individual assay of both isoenzymes, one abundant in  
muscle, of pyruvate kinase (PK), recently reported superior to serum  
creatine phosphokinase (CPK) in detecting patients with and female carriers  
of X-linked recessive (Duchenne) muscular dystrophy (DMD). Extensive  
comparative studies did not support these findings and confirmed the marked  
superiority of CPK over variants of PK or other enzymes in sensitivity,  
stability and convenience. Deducting the adenylate kinase increment (AKI)  
further refined the CPK assay, eliminating the effect of **hemolysis** in  
diagnosis and enabling studies of blood cell content. Both leucocytes and  
erythrocytes liberated PK and lactate dehydrogenase (LDH) after brief  
chilling or disruption. Only erythrocytes showed a CPK content, however,  
constantly adjusted to match that of serum as if by free cell membrane  
passage, but less accomodating to a sudden large influx of CPK than of LDH,  
where an apparent buffering effect could account for differences in  
clinical response.  
? logoff y

```

06aug98 16:51:43 User208670 Session D657.4
    $0.74      0.247 DialUnits File155
        $3.40  17 Type(s) in Format  7
        $3.40  17 Types
$4.14 Estimated cost File155
    $0.38      0.072 DialUnits File5
        $7.25  5 Type(s) in Format  7
        $7.25  5 Types
$7.63 Estimated cost File5
    $0.76      0.098 DialUnits File73
        $11.40 6 Type(s) in Format  7
        $11.40 6 Types
$12.16 Estimated cost File73
    $0.15      0.046 DialUnits File144
$0.15 Estimated cost File144
    $0.07      0.015 DialUnits File434
$0.07 Estimated cost File434
    $0.20      0.046 DialUnits File34
$0.20 Estimated cost File34
    $0.15      0.031 DialUnits File76
$0.15 Estimated cost File76
    $0.11      0.015 DialUnits File305
$0.11 Estimated cost File305
    $0.06      0.026 DialUnits File156
$0.06 Estimated cost File156
    $0.11      0.026 DialUnits File442
$0.11 Estimated cost File442
    $0.08      0.021 DialUnits File35
$0.08 Estimated cost File35
    $0.10      0.036 DialUnits File50
$0.10 Estimated cost File50
    $0.07      0.026 DialUnits File77
$0.07 Estimated cost File77
    $0.11      0.021 DialUnits File150
$0.11 Estimated cost File150
    $0.08      0.021 DialUnits File376
$0.08 Estimated cost File376
    $0.14      0.026 DialUnits File2
$0.14 Estimated cost File2
    $0.11      0.021 DialUnits File8
$0.11 Estimated cost File8
    $0.10      0.021 DialUnits File103
$0.10 Estimated cost File103
    $0.08      0.026 DialUnits File348
$0.08 Estimated cost File348
    $0.44      0.088 DialUnits File653
        $0.60  1 Type(s) in Format  3 (UDF)
        $0.60  1 Types
$1.04 Estimated cost File653
    $0.36      0.072 DialUnits File654
$0.36 Estimated cost File654
    OneSearch, 21 files, 1.000 DialUnits FileOS
$27.05 Estimated cost this search
$27.05 Estimated total session cost 1.000 DialUnits
Logoff: level 98.07.06 D 16:51:44

```